

Amendments to the Claims

The listing of claims will replace all prior versions, and listings of claims in the application.

1. (Previously presented) An isolated polypeptide of the structure or formula S-(L)_n-B wherein:

- (a) S is selected from the group consisting of PTH(1-9) (AlaValSerGluIleGlnLeuMetHis) (SEQ ID NO: 1), PTH(1-5) (AlaValSerGluIle) (SEQ ID NO: 4) or PTH (1-11) (AlaValSerGluIleGlnLeuMetHisAsnLeu) (SEQ ID NO: 46);
- (b) L is a glycine present n times;
- (c) n is 5 to 10; and
- (d) B is a carboxy terminal binding domain of PTH(1-34) or PTHrP(1-34), wherein said carboxy terminal binding domain binds to a PTH- receptor 1 molecule wherein said polypeptide stimulates intracellular accumulation of cyclic cAMP.

2-4. (Canceled)

5. (Previously presented) The isolated polypeptide of claim 1, wherein L is selected from the group consisting of Gly₅, Gly₇ and Gly₉.

6. (Previously presented) The isolated polypeptide of claim 1, wherein B is selected from the group consisting of PTH(15-31)

(LeuAsnSerMetGluArgValGluTrpLeuArgLysLysLeuGlnAspVal) (SEQ ID NO:2),
PTH(17-31) (SerMetGluArgValGluTrpLeuArgLysLysLeuGlnAspVal) (SEQ ID NO:63),
PTHrP (15-31) (IleGlnAspLeuArgArgArgPhePheLeuHisHisLeuIleAlaGluIle) (SEQ ID
NO:8), and PTHrP(17-31) (AspLeuArgArgArgPhePheLeuHisHisLeuIleAlaGluIle) (SEQ
ID NO:12).

7. (Previously presented) An isolated polypeptide selected from the group
consisting of: PG5: AlaValSerGluIleGlnLeuMetHisGlyGlyGlyGlyGlyLeuAsnSer
MetGluArgValGluTrpLeuArgLysLysLeuGlnAspVal (SEQ ID NO:3), PG9:
AlaValSerGluIleGlyGlyGlyGlyGlyGlyGlyGlyGlyGlyLeuAsnSerMetGluArgValGluTrp
LeuArgLysLysLeuGlnAspVal (SEQ ID NO:5), PG7: AlaValSerGluIleGlnLeu
MetHisGlyGlyGlyGlyGlyGlyGlySerMetGluArgValGluTrpLeuArgLysLysLeuGlnAsp
Val (SEQ ID NO:6), PrPG5:AlaValSerGluHisGlnLeuLeuHisGlyGlyGlyGlyGlyIle
GlnAspLeuArgArgArgPhePheLeuHisHisLeuIleAlaGluIle (SEQ ID NO:64), PrPG9:
AlaValSerGluHisGlyGlyGlyGlyGlyGlyGlyGlyGlyGlyIleGlnAspLeuArgArgArg
PhePheLeuHisHisLeuIleAlaGluIle (SEQ ID NO:65) and PrPG7: AlaValSerGlu
HisGlnLeuLeuHisGlyGlyGlyGlyGlyGlyGlyAspLeuArgArgArgPhePheLeuHisHisLeuIle
AlaGluIle (SEQ ID NO:66).

8. (Previously presented) An isolated polypeptide selected from the group
consisting of PG5: AlaValSerGluIleGlnLeuMetHisGlyGlyGlyGlyGlyLeuAsnSer
MetGluArgValGluTrpLeuArgLysLysLeuGlnAspVal (SEQ ID NO:3), PG9:
AlaValSerGluIleGlyGlyGlyGlyGlyGlyGlyGlyGlyGlyLeuAsnSerMetGluArgVal

GluTrpLeuArgLysLysLeuGlnAspVal (SEQ ID NO:5), PG7: AlaValSerGluIleGlnLeu
MetHisGlyGlyGlyGlyGlyGlyGlySerMetGluArgValGluTrpLeuArgLysLysLeuGlnAsn
Val (SEQ ID NO:6).

9. (Canceled)

10. (Previously presented) An isolated polypeptide of the structure or formula S-
(L)_n-B wherein:

- (a) S is X Val X Glu X X X X His (SEQ ID NO: 42), wherein X is an amino acid;
- (b) L is glycine and n equals 5-10; and
- (c) B is a carboxy terminal binding domain of PTH(1-34) or PTHrP(1-34), wherein said carboxy terminal binding domain binds to a PTH-receptor 1 molecule, wherein said polypeptide stimulates intracellular accumulation of cyclic cAMP.

11. (Previously presented) An isolated polypeptide of the structure or formula S-
(L)_n-B wherein :

- (a) S is Ser Val Ser Glu Ile Gln Leu Met His (SEQ ID NO: 44);
- (b) L is 5-10 glycine residues; and
- (c) B is Leu Asn Ser Met Glu Arg Val Glu Trp Leu Arg Lys Lys Leu Gln Asp Val (SEQ ID NO: 45).

12-13. (Canceled).

14. (Previously presented) An isolated polypeptide encoded by a nucleic acid sequence selected from the group consisting of: SEQ ID NO:14, SEQ ID NO:15 and SEQ ID NO:16.

15-43. (Canceled)

44. (Previously presented) The isolated polypeptide of claim 7, wherein said polypeptide is modified to improve the solubility, absorption, or biological half-life of said polypeptide and wherein said modification is selected from the group consisting of the addition of C₁₋₁₂ alkyl groups, the addition of C₁₋₁₂ hydroxyalkyl groups, the addition of acyl groups, and lactam cyclization.

45. (Previously presented) The isolated polypeptide of claim 8, wherein said polypeptide is modified to improve the solubility, absorption, or biological half-life of said polypeptide and wherein said modification is selected from the group consisting of the addition of C₁₋₁₂ alkyl groups, the addition of C₁₋₁₂ hydroxyalkyl groups, the addition of acyl groups, and lactam cyclization.

46. (Previously presented) The isolated polypeptide of claim 1, wherein n is an integer from 5 to 9.